

(4)

FILE COPY

MEAN SQUARE ERROR BEHAVIOR FOR PREDICTION  
IN LINEAR REGRESSION MODELS

BY

ALAN E. GELFAND

TECHNICAL REPORT NO. 427

MARCH 7, 1990

PREPARED UNDER CONTRACT

N00014-89-J-1627 (NR-042-267)

FOR THE OFFICE OF NAVAL RESEARCH

AD-A219 988

Reproduction in Whole or in Part is Permitted  
for any purpose of the United States Government

Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA



DTIC  
ELECTE  
APR 03 1990

cc E

D

90 04 02 121

MEAN SQUARE ERROR BEHAVIOR FOR PREDICTION  
IN LINEAR REGRESSION MODELS

BY

ALAN E. GELFAND

TECHNICAL REPORT NO. 427

MARCH 7, 1990

Prepared Under Contract

N00014-89-J-1627 (NR-042-267)

For the Office of Naval Research

Herbert Solomon, Project Director

Reproduction in Whole or in Part is Permitted  
for any purpose of the United States Government



Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

# MEAN SQUARE ERROR BEHAVIOR FOR PREDICTION IN LINEAR REGRESSION MODELS

Alan E. Gelfand

## ABSTRACT

For the problem of individual prediction in linear regression models, that is, estimation of a linear combination of regression coefficients, mean square error behavior of a general class of adaptive predictors is examined. jha

## 1. INTRODUCTION

Suppose the usual linear regression model with fixed regressors,  $Y = X\beta + \epsilon$ ,  $Y_{n \times 1}$ ,  $X_{n \times p}$  full rank,  $\beta_{p \times 1}$  and  $\epsilon_{n \times 1} \sim (0, \sigma^2 I)$ . Let  $\hat{\beta}_{LS} = (X^T X)^{-1} X^T Y$  denote the ordinary least squares estimator of  $\beta$ . At a new vector of predictor values,  $X_0$ , we seek to estimate  $X_0^T \beta$ . Using mean square error as a criterion, results of Cohen (1965) show that if  $\epsilon$  is normally distributed,  $\alpha X_0^T \hat{\beta}_{LS}$  is an admissible estimator of  $X_0^T \beta$  for  $0 \leq \alpha \leq 1$ , e.g., the UMVU predictor is admissible. In fact, a predictor of the form  $l^T Y$  is admissible for  $X_0^T \beta$  iff  $(2I - X(X^T X)^{-1} X^T) X_0 \leq X_0^T (X^T X)^{-1} X_0$ .

In the sequel, we study the MSE under normality of predictors of the form  $X_0^T \hat{\beta}_C$  where

$$\hat{\beta}_C = C \hat{\beta}_{LS} + (I - C) \beta^* \quad (1)$$

$C$  a matrix usually data dependent and  $\beta^*$  a specified vector. Such  $\hat{\beta}_C$  include most alternatives to  $\hat{\beta}_{LS}$  discussed in the literature. Earlier work in this direction appears in Baranchik (1964) and Radhakrishnan (1970).

## 2. NOTATION AND MOTIVATION

To simplify matters, we convert to canonical form. Let  $\hat{\alpha} = P\hat{\beta}_{LS}$ ,  $P$  orthogonal such that  $P(X^T X)^{-1}P^T = D^{-1}$ ,  $D$  diagonal with diagonal elements  $d_i$ . Define  $\alpha = P\beta$ ,  $\ell = PX_0$  and for convenience set  $\beta^* = 0$ . For the moment assume  $\sigma^2$  known. Our problem now is to estimate  $\theta = \ell^T \alpha$  given  $\hat{\alpha} \sim N(\alpha, \sigma^2 D^{-1})$  wishing to do well near  $\theta = 0$ . Let  $U = \ell^T \hat{\alpha}$ ,  $Z = \hat{\alpha}^T D \hat{\alpha}$ ,  $q = \ell^T D^{-1} \ell$ ,  $V = Z - U^2/q$ ,  $\lambda = \alpha^T D \alpha$  and  $\zeta = \lambda - \theta^2/q$ . Then,  $U, V$  are independent,  $U \sim K(\theta, \sigma^2 q)$ ,  $V \sim \sigma^2 \chi_{p-1}^2(\zeta/\sigma^2)$ .

Consider a general adaptive predictor  $\delta(\hat{\alpha})$  of the form

$$\delta(\hat{\alpha}) = \sum h_i(\hat{\alpha}) \ell_i \hat{\alpha}_i. \quad (2)$$

Most predictors of  $\theta$  discussed in the literature are special cases of (2). Apart from the LS predictor,  $U$ , we have:

i) A class of predictors given in Thompson (1968)

$$T_m = \frac{U^2}{U^2 + m\sigma^2 q} U, \quad m \text{ a known constant, i.e., } h_i(\hat{\alpha}) = \frac{(\hat{\alpha}^T \ell)^2}{(\hat{\alpha}^T \ell)^2 + m\sigma^2 q}.$$

ii) A class of predictors given in Mehta and Srivastava (1971)

$$MS_{b_1, b_2} = (1 - b_1 e^{-b_2 U^2 / \sigma^2 q}) U, \quad 0 < b_1 < 1, \quad b_2 > 0, \quad b_1, b_2 \text{ known,}$$

$$\text{i.e., } h_i(\hat{\alpha}) = 1 - b_1 \exp(-b_2 (\hat{\alpha}^T \ell)^2 / \sigma^2 q).$$

iii) A predictor arising from the James-Stein estimator adapted for unequal variances (Sclove 1968)

$$JS_c = (1 - \frac{c\sigma^2}{2}) U, \quad c \text{ known usually taken equal to } p - 2.$$

A positive part adjustment should be applied so that  $h_i(\hat{\alpha}) = [1 - c\sigma^2 (\hat{\alpha}^T D \hat{\alpha})^{-1}]^+.$

iv) Predictors arising from (simple) ridge estimators

$$R_{k_t} = \sum_i \frac{d_i}{d_i + k_t} \hat{\alpha}_i$$

where  $k_t$  is based on the data, i.e.,  $h_i(\hat{\alpha}) = d_i / (d_i + k_t(\hat{\alpha}))$ .

$k$ 's discussed include:

$$k_1(\hat{\alpha}) = \sigma^2 p (\hat{\alpha}^T \hat{\alpha})^{-1} \quad (\text{Hoerl, Kennard, and Baldwin 1975}),$$

$$k_2(\hat{\alpha}) = \sigma^2 p Z^{-1} \quad (\text{Lawless and Wang 1976}),$$

$$k_3(\hat{\alpha}), \text{ the solution to } \sum_i \hat{\alpha}_i^2 d_i^2 (d_i + k_3)^{-2} = \sum_i \hat{\alpha}_i^2 - \sigma^2 \sum_i d_i^{-1}$$

(McDonald and Galarneau 1975),

$$k_4(\hat{\alpha}), \text{ the solution to } \sum_i \hat{\alpha}_i^2 d_i (d_i + k_4)^{-1} = \sigma^2 p$$

(the RIDGM estimator of Dempster, Schatzoff and Wermuth 1977).

A subclass of (2) which includes (i), (ii), (iii), and  $R_{k_2}$  has the form

$$\delta(\hat{\alpha}) = \sum_i h_i(U, Z) \hat{\alpha}_i \quad (3)$$

A further subclass which still includes (i), (ii), and (iii) is

$$\delta(\hat{\alpha}) = h(U, Z) \cdot U. \quad (4)$$

When  $D = I$ , all of the aforementioned estimators belong to (4).

Taking another point of view (see e.g. Thompson (1968)), if  $h_i$  in (3) is constant, the optimal  $h_i$  to minimize the MSE are easily obtained:

$$h_i^* = \frac{6}{\sigma^2 + \lambda} \frac{\alpha_i}{\ell_i} \quad (5)$$

An estimator of  $h_i^*$  would be of the form  $c_i(\hat{\alpha}, \sigma^2)$  leading to a predictor belonging to (2). If (5) was estimated by  $c(U, Z, \sigma^2) \cdot \hat{\alpha}_i / \ell_i$  the class (4) results.

Suppose we take a Bayesian approach using a prior which centers  $\theta$  at 0, where we want to do well. More precisely, let  $Q$  be an orthogonal matrix such that  $QD^{\frac{1}{2}}\alpha = \begin{pmatrix} \epsilon/\sqrt{q} \\ \eta \end{pmatrix}$  where  $\eta$  is  $(p-1) \times 1$  and  $\eta^T \eta = \phi$ . If we take as our prior

$$(\theta/\sqrt{q}) \sim N(0, \begin{pmatrix} \gamma & 0 \\ 0 & \rho \gamma I_{p-1} \end{pmatrix}), \rho \text{ known,}$$

then under squared error loss, the Bayes estimate of  $\theta$  is  $(\gamma + \sigma^2)^{-1} \cdot \gamma U$ . Since  $(U, Z)$  is sufficient under the marginal distribution of  $\omega = QD^{\frac{1}{2}}\hat{\alpha}$  an "empirical Bayes" estimator of  $\theta$  takes the form in (4).

### 3. EXAMINATION OF THE MSE

We can calculate the MSE for the general predictor in (2) in terms of the  $h_i$ , assuming  $\sigma^2$  known.<sup>1</sup>

Theorem 1. If  $E|\frac{\partial h_i}{\partial U} \cdot \hat{\alpha}_i| < \infty$ ,  $i = 1, 2, \dots, p$ ,

$$\begin{aligned} \text{MSE}(\delta) &= \sigma^2 q + E(\delta - U)^2 - 2\sigma^2 E \sum_i \hat{\alpha}_i^2 (1 - h_i) \\ &\quad + 2\sigma^2 q E \sum_i \hat{\alpha}_i \frac{\partial h_i}{\partial U}. \end{aligned} \quad (6)$$

Proof. By direct calculation

$$\text{MSE}(\delta) = \sigma^2 q + E(\delta - U)^2 - 2E\{r(\hat{\alpha})(U - \delta)\} \quad (7)$$

where  $r(\hat{\alpha}) = \sum (1 - h_i) \hat{\alpha}_i$ . Stein's identity (Stein 1981, p. 1148) converts the right-most term of (7) to  $\sigma^2 q E(\frac{\partial r(\hat{\alpha})}{\partial U})$ . Simplification yields (6).

$\frac{\partial h_i}{\partial U}$  would be calculated using the transformation  $\hat{\alpha} = D^{-\frac{1}{2}} Q^T u$  of the previous section. In the case of (3), it can be calculated directly writing  $h_i$  as a function of  $U$  and  $V$ . For predictors of the form (4),  $\text{MSE}(\delta)$  depends only on  $\theta$  and  $\phi$  and is given as

Corollary 1.

Corollary 1. For the predictors in (4), if  $E|U \frac{\partial h}{\partial U}| < \infty$

$$\text{MSE}(\delta) = \sigma^2 q + E(1 - h)^2 U^2 + 2\sigma^2 q E U \frac{\partial h}{\partial U} - 2\sigma^2 q E(1 - h). \quad (8)$$

Under (4) choices of  $h$  in the literature are such that  $h$  is symmetric in  $U$  about 0 and restricted to  $[0, 1]$ . Using essentially

the argument of Efron and Morris (1976, p. 14) positive part restriction of  $h$  uniformly reduces risk. Restriction of  $h \leq 1$  is less clear. Taking  $h > 0$  the predictor  $h^* \cdot U$  where  $h^* = \min(h, 1)$  does not necessarily dominate  $h \cdot U$ . For example, let

$$h(U, V) = \begin{cases} 1 + c, & a^2 < U^2 < b^2 \\ 1, & \text{elsewhere} \end{cases}.$$

Then at each  $\phi$ , for  $|\theta|$  sufficiently large, MSE of  $h(U, V)U$  is less than MSE of  $h^*(U, V)$ . Nonetheless, to improve in a neighborhood of a specified  $\theta_0$  requires

convex combinations of  $U$  and  $\theta_0$ . Theorem 2 details MSE properties of predictors in (4) relative to the MSE of  $U$ .

Theorem 2. For  $\delta(\hat{\alpha})$  in (4) with  $h \in [0, 1]$ , let  $h$  be symmetric in  $U$  about 0. Let  $g = (1 - h)U$  with  $\limsup_{|U| \rightarrow \infty} g = 0$  and assume  $\frac{\partial g}{\partial U}$

exists for all  $U$ . Finally, assume that the Lebesgue measure of  $A = \{(U, V) : h(U, V) < 1\}$  is greater than 0. Then,

(i) For each  $\phi$  there is a neighborhood  $N_\phi$  of  $\theta = 0$  where  $\text{MSE}(\delta; \theta, \phi) < \sigma^2 q$ .

(ii)  $\text{MSE}(\delta; \theta, \phi)$  is bounded and  $\lim_{|\theta| \rightarrow \infty} \text{MSE}(\delta; \theta, \phi) = \sigma^2 q$ .

(iii)  $\text{MSE}(\delta; \theta, \phi)$  is symmetric in  $\theta$  about 0 and  $\left. \frac{\partial \text{MSE}(\delta; \theta, \phi)}{\partial \theta} \right|_{\theta=0} = 0$ .

(iv)  $g^2 - 2 \frac{\partial g}{\partial U}$  changes sign at least once in  $0 < U < \infty$ . If  $g^2 - 2 \frac{\partial g}{\partial U}$  changes sign  $b$  times in  $0 < U < \infty$ , then for fixed  $\phi$ ,  $\text{MSE}(\delta; \theta, \phi) - \sigma^2 q$  changes sign at most  $2b$  times.

Proof. The proof of (i) is clear since  $\text{MSE}(\delta; 0, \phi) < \sigma^2 q$ . For (ii),

$$\text{MSE}(\delta; \theta, \phi) = \sigma^2 q + E g^2 - 2E(U - \theta)g. \quad (9)$$

Given  $\epsilon$ ,  $\exists u_0$  such that for all  $V$ ,  $U > u_0 \Rightarrow |g| < \epsilon$  and  $\exists \theta_0 > 0$  such that  $|\theta| > \theta_0 \Rightarrow P(|U| > u_0) > 1 - \epsilon$ . Then the second term and the third term (using the Cauchy-Schwarz Inequality) in (9) can be made arbitrarily small as  $|\theta| \rightarrow \infty$ . It is clear that the r.h.s. of (9) is bounded. (iii) is obvious. The first part of (iv) follows since  $U$  is admissible. The second part follows from

the sign change theorem of Karlin (1957) by noting that  $MSE(\delta; \epsilon, \phi) - \sigma^2 q = E(g^2 - 2 \frac{\partial g}{\partial U})$ .

Remark 1. Predictors in (i), (ii), (iii) of Section 2 satisfy the conditions of Theorem 2.

Remark 2. Result (ii) is a simple case of the "tail minimaxity" notion of Berger (1976).


Remark 3. In (iii),  $\inf_{\epsilon} MSE(\delta; \epsilon, \phi)$  need not occur at  $\epsilon = 0$ . If, however,  $h(U, V)$  is increasing in  $|U|$  it must as may be shown by establishing the result for  $h$ , a step function in  $U$ . An induction argument proves this.

Remark 4. If  $b = 1$  in (iv), then a graph of  $MSE(\delta; \epsilon, \phi)$  for  $\epsilon \geq 0$  must start below  $c^2 q$  at  $\epsilon = 0$ , cross above  $c^2 q$  at some  $\epsilon$  and then asymptotically return to  $\sigma^2 q$  from above. Any predictor satisfying the conditions of Theorem 2 must necessarily perform worse for a set of  $\epsilon$ 's near 0 than for a set arbitrarily far away.

Remark 5. No immediate extension of Theorem 2 to  $\delta(\hat{q})$  as in (3) is available. For an arbitrary member of (3),  $MSE$  depends upon  $\theta$  and  $n$  and, even if each  $h_i$  meets the "tail minimaxity" condition, need not approach  $\sigma^2 q$  as  $|\theta| \rightarrow \infty$  for fixed  $n$ .

Remark 6. Theorem 2 is readily extended to the comparison of any pair of predictors in (4).

We conclude with a comment on admissibility for the above predictors. Within the class of predictors based solely on  $U$ , i.e.,  $h(U)U$ , those meeting the conditions of Theorem 2 will either be admissible or if not then improvement cannot be substantial. We employ ideas of Chow and Hwang (1984). Suppose  $\delta_1(U)$  is to dominate  $\delta_0 = h(U)U$  meeting the conditions of Theorem 2. We can write  $\delta_1$  as  $h^*(U)U$ , and assume  $h^* \geq 0$ . For  $\delta_1$  to dominate  $\delta_0$  requires, when  $|U|$  is large, that generally  $h^*$  be closer to 1 than  $h$  and that, when  $|U|$  is small, generally  $h^*$  be closer to 0 than  $h$ . A simplified picture of  $\delta_0, \delta_1$  for  $U > 0$  might look like



But, at  $\epsilon = \epsilon_0$ , it would be almost impossible for  $\epsilon_1$  to dominate. Thus, the simplest  $h^*$  which realistically could dominate would have to have at least 3 sign changes for  $h - h^*$  on  $U > 0$ . For such an  $h^*$ , its form would be complicated, domination would be difficult to show, and improvement would be minimal.

This argument does not extend to the more general class (4). Though  $U$  and  $V$  are independent, conditioning on  $V$  in the above heuristic leads to  $\epsilon_0$  depending upon  $V$ . We, nonetheless, conjecture "approximate admissibility" for members of (4) meeting the conditions of Theorem 2.

#### FOOTNOTE

<sup>1</sup>When  $\sigma^2$  is unknown, we customarily assume an estimator  $S^2$  of  $\sigma^2$  such that  $vS^2 \sim c^2\chi_v^2$  independent of  $\hat{\alpha}$ . In the foregoing predictors,  $\sigma^2$  is replaced by  $cS^2$ . As Lawless (1981, pp. 463-464) notes, when  $v \rightarrow \infty$  and even when  $v$  is moderate, resulting MSE will differ little from that with  $\sigma^2$  known.

#### BIBLIOGRAPHY

- Baranchik, Alvin J. (1964). "Multiple Regression and Estimation of the Mean of a Multivariate Normal Distribution," Stanford University Tech. Report #51.
- Berger, J.O. (1976). "Tail minimaxity in location vector problems and its applications," Annals of Statistics, 4, 33-50.
- Chow, M.S. and Hwang, J.T. (1984). "The comparison of estimators for the noncentrality of a chi-square distribution," Cornell Statistical Center, Technical Report.
- Cohen, A. (1965). "Estimates of linear combinations of the parameters in the mean vector of a multivariate distribution," Annals of Mathematical Statistics, 36, 78-87.
- Dempster, A.P., Scharzoff, M. and Wermuth, N. (1977). "A simulation study of alternatives to ordinary least squares," Journal of the American Statistical Association, 72, 77-106.
- Efron, B. and Morris, C. (1976). "Families of minimax estimators of the mean of a multivariate normal distribution," Annals of Statistics, 4, No. 1, 11-21.
- Hoerl, A.E., Kennard, R.W. and Baldwin, K.F. (1975). "Ridge regression: Some simulations," Communications in Statistics, 4, 105-123.
- Karlin, S. (1957). "Polya type distributions II," Annals of Mathematical Statistics, 28, No. 2, 281-308.
- Lawless, J.F. (1981). "Mean squared error properties of generalized ridge estimators," Journal of the American Statistical Association, 76, 462-466.
- Lawless, J.F. and Wang, P. (1976). "A simulation study of ridge and other regression estimators," Communications in Statistics, A, 5, 307-323.
- McDonald, G.C. and Galarneau, D.L. (1975). "A Monte Carlo evaluation of some ridge type estimators," Journal of the American Statistical Association, 70, 407-416.
- Mehta, J.S. and Srivastava, R. (1971). "Estimation of the mean by shrinking to a point," Journal of the American Statistical Association, 66, 86-91.

- Radhakrishnan, R. (1970). "Some Estimation Problems in Multivariate Analysis," unpublished Ph.D. Thesis, Carnegie Mellon University.
- Sclove, S.L. (1968). "Improved estimators for coefficients in linear regression," Journal of the American Statistical Association, 63, 596-606.
- Stein, C. (1981). "Estimation of the mean of a multivariate normal distribution," Annals of Statistics, 9, No. 6, 1135-1151.
- Thompson, J.R. (1968). "Some shrinkage techniques for estimating the mean," Journal of the American Statistical Association, 63, 113-122.

**UNCLASSIFIED**

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER  427	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  Mean Square Error Behavior For Prediction In Linear Regression Models		5. TYPE OF REPORT & PERIOD COVERED  TECHNICAL REPORT
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  Alan E. Gelfand		8. CONTRACT OR GRANT NUMBER(s)  N00014-89-J-1627
9. PERFORMING ORGANIZATION NAME AND ADDRESS  Department of Statistics Stanford University Stanford, CA 94305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  NR-042-267
11. CONTROLLING OFFICE NAME AND ADDRESS  Office of Naval Research Statistics & Probability Program Code 1111		12. REPORT DATE  March 7, 1990
		13. NUMBER OF PAGES  11
		14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)
		15. SECURITY CLASS (of this report)  UNCLASSIFIED
		16a. DECLASSIFICATION/DOWNGRADING SCHEDULE
18. DISTRIBUTION STATEMENT (of this Report)  APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
19. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  prediction; adaptive estimators; linear regression; mean square error.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  For the problem of individual prediction in linear regression models, that is, estimation of a linear combination of regression coefficients, mean square error behavior of a general class of adaptive predictors is examined.		